

PATH INTEGRAL ON RELATIVISTIC SPINLESS POTENTIAL PROBLEMS

De-Hone Lin

Department of Physics, National Tsing Hua University

Hsinchu 30043, Taiwan

E-mail: d793314@phys.nthu.edu.tw

The formulation of the relativistic spinless path integral on the general affine space is presented. For the one dimensional space, the Duru-Kleinert (DK) method and the δ -function perturbation technique are applied to solve the relativistic path integrals of the smooth potential and the Dirichlet boundary condition problems, respectively.

1 Relativistic Path Integral on General Affine Space

Let us first consider a point particle of mass M moving at a relativistic velocity in a $(D + 1)$ -dimensional Minkowski space with a given electromagnetic field. By using $t = -i\tau = -ix^4$, the path integral representation of the fixed-energy amplitude is conveniently formulated in a $(D + 1)$ -Euclidean spacetime with the Euclidean metric,

$$(g_{\mu\nu}) = \text{diag} (1, \dots, 1, c^2), \quad (1)$$

and is given by^{1,3}

$$G(\mathbf{x}_b, \mathbf{x}_a; E) = \frac{i\hbar}{2Mc} \int_0^\infty dL \int \mathcal{D}\rho(\lambda) \Phi[\rho(\lambda)] \int \mathcal{D}^D x(\lambda) e^{-A_E/\hbar} \quad (2)$$

with the action

$$A_E = \int_{\lambda_a}^{\lambda_b} d\lambda \left[\frac{M}{2\rho(\lambda)} \mathbf{x}'^2(\lambda) - i\frac{e}{c} \mathbf{A}(\mathbf{x}) \cdot \mathbf{x}'(\lambda) - \rho(\lambda) \frac{(E - V(\mathbf{x}))^2}{2Mc^2} + \rho(\lambda) \frac{Mc^2}{2} \right], \quad (3)$$

where L is defined by

$$L = \int_{\lambda_a}^{\lambda_b} d\lambda \rho(\lambda), \quad (4)$$

in which $\rho(\lambda)$ is an arbitrary dimensionless fluctuating scale variable, and $\Phi[\rho]$ is some convenient gauge-fixing functional, such as $\Phi[\rho] = \delta[\rho - 1]$, to fix the value of $\rho(\lambda)$ to unity. $\hbar/Mc$ is the well-known Compton wave length of a particle of mass M , $\mathbf{A}(\mathbf{x})$ is the vector potential, $V(\mathbf{x})$ is the scalar potential,

E is the system energy, and $\mathbf{x}$ is the spatial part of the $(D+1)$ vector $x = (\mathbf{x}, \tau)$. This path integral forms the basis for studying relativistic potential problems. To obtain a tractable path integral in (2) for the arbitrary time independent potential $V(\mathbf{x})$, Let us first take the space-time transformation as usual²

$$\Delta\lambda_n = \Delta s_n f(\mathbf{x}_n), \quad (5)$$

where $\Delta\lambda_n \equiv \lambda_n - \lambda_{n-1}$ is a small parameter interval and $f(\mathbf{x})$ are invertible but otherwise arbitrary function. A general formulation of relativistic path integral in affine space is given by introducing the coordinate transformation

$$x^i = h^i(q). \quad (6)$$

The differential mapping may be written as

$$dx^i = \partial_\mu h^i(q) dq^\mu = e^i{}_\mu(q) dq^\mu. \quad (7)$$

Since we must find all terms that will eventually contribute to order ϵ , we expand $(\Delta\mathbf{x}_n)^2 = (\mathbf{x}_n - \mathbf{x}_{n-1})^2$ up to fourth order in $\Delta q_n^\mu = q_n^\mu - q_{n-1}^\mu$. Finally, we obtain the path integral representation for the fixed-energy amplitude of a relativistic particle in general affine space^{1,2,3}

$$\begin{aligned} G(\mathbf{x}_b, \mathbf{x}_a; E) = & \lim_{n \rightarrow \infty} \frac{i\hbar}{2Mc} \int_0^\infty dS \frac{f(q_a)}{\left(\frac{2\pi\hbar\epsilon_n^s \rho_b f(q_a)}{M}\right)^{D/2}} \prod_{n=1}^{N+1} \left[\int d\rho_n \Phi(\rho_n) \right] \\ & \times \prod_{n=2}^{N+1} \left[\int_{-\infty}^\infty \frac{d^D \Delta q_n g^{1/2}(q_n)}{\left(\frac{2\pi\hbar\epsilon_n^s \rho_n f(q_n)}{M}\right)^{D/2}} \right] \exp \left\{ -\frac{1}{\hbar} \sum_{n=1}^{N+1} [A^\epsilon + A_J^\epsilon + A_{\text{pot}}^\epsilon] \right\} \end{aligned} \quad (8)$$

with each s -sliced action

$$\begin{aligned} A^\epsilon + A_J^\epsilon + A_{\text{pot}}^\epsilon = & \frac{M}{2\epsilon^s \rho f} g_{\mu\nu} \Delta q^\mu \Delta q^\nu - \frac{\hbar}{2} \Gamma_\mu{}^\mu{}_\nu \Delta q^\nu + \epsilon^s \rho f \frac{\hbar^2}{8M} (\Gamma_\mu{}^\mu{}_\nu)^2 \\ & - i \frac{e}{c} \left[A_\mu \Delta q^\mu - \epsilon^s \rho f \frac{\hbar}{2M} (A_\nu \Gamma_\mu{}^{\mu\nu} + D_\mu A^\mu) \right] - \epsilon^s \rho f \frac{(E - V(q))^2}{2Mc^2}, \end{aligned} \quad (9)$$

where we have omitted the subscripts n of Δq^μ . The induced metric $g_{\mu\nu} = e^i{}_\mu e^i{}_\nu$ and affine connection $\Gamma_{\lambda\kappa}{}^\mu = e_i{}^\mu e^i{}_{\kappa,\lambda}$ are evaluated at the postpoint, $\dot{q}^\alpha$ stands for dq^α/ds . The action of covariant derivative D_μ is defined as

$$D_\mu A^\mu = \partial_\mu A^\mu + \Gamma_\mu{}^\mu{}_\lambda A^\lambda. \quad (10)$$

Remarkably, this expression involves only the connection contracted in the first two indices:

$$\Gamma_\mu{}^{\mu\nu} = g^{\mu\lambda} \Gamma_{\mu\lambda}{}^\nu. \quad (11)$$

2 Duru-Kleinert Method for the Relativistic Potential Problems

The relativistic stable path integrals of Eq. (2) have a more elegant representation if the systems are in two-dimensional Minkowski space or rotationally invariant systems in any dimensions. By decomposing the Eq. (2) into angular parts and take DK transformation, we get ²

$$G(\mathbf{x}_b, \mathbf{x}_a; E) = \frac{1}{(r_b r_a)^{(D-1)/2}} \sum_{l=0}^{\infty} G_l^{\text{DK}}(r_b, r_a; E) \sum_{\mathbf{m}} Y_{l\mathbf{m}}(\hat{\mathbf{x}}_b) Y_{l\mathbf{m}}^*(\hat{\mathbf{x}}_a), \quad (12)$$

where superscript DK indicates that the system has been performed by the DK-transformation, the functions $Y_{l\mathbf{m}}(\hat{\mathbf{x}})$ are the D -dimensional hyperspherical harmonics and $G_l^{\text{DK}}(r_b, r_a; E)$ is the purely radial transformed amplitude

$$G_l^{\text{DK}}(r_b, r_a; E) = \frac{\hbar i}{2Mc} f_b^{1/4} f_a^{1/4} G(q_b, q_a; E). \quad (13)$$

The amplitude $G(q_b, q_a; E)$ is given by ²

$$G(q_b, q_a; E) \equiv \int_0^\infty dS \int \mathcal{D}\rho(s) \Phi[\rho(s)] \int \mathcal{D}q(s) e^{-A_s^{\text{DK}}/\hbar} \quad (14)$$

with

$$A_s^{\text{DK}} = \int_0^S ds \left[\frac{M}{2\rho(s)} \dot{q}^2(s) + \rho(s) f(q(s)) \right. \\ \left. \times \left(\frac{\hbar^2}{2M} \frac{(l + D/2 - 1)^2 - 1/4}{r^2(q(s))} - \frac{[E - V(r(q))]}{2Mc^2} + \frac{Mc^2}{2} \right) + V_{\text{eff}} \right]. \quad (15)$$

The effective potential V_{eff} is given by

$$V_{\text{eff}} = -\frac{\rho(s)\hbar^2}{M} \left[\frac{1}{4} \frac{h'''(q)}{h'(q)} - \frac{3}{8} \left(\frac{h''(q)}{h'(q)} \right)^2 \right]. \quad (16)$$

Here the transformation function $r = h(q)$ with

$$h'^2(q) = f(r). \quad (17)$$

3 Relativistic Path Integral With Dirichlet Boundary Conditions via Sum Over Perturbation Series

With the help of the δ -function perturbation techniques, we obtain the relativistic fixed-energy amplitude for an impenetrable wall appears at $x = a$ given

by³

$$G^{(\text{wall})}(x_b, x_a; E) = G_0(x_b, x_a; E) - \frac{G_0(x_b, a; E)G_0(a, x_a; E)}{G_0(a, a; E)}. \quad (18)$$

Furthermore, for the relativistic motion inside a box with boundaries at $x = a$ and $x = b$ one obtains ($a < x < b$, Dirichlet-Dirichlet boundary conditions)³

$$G(x_b, x_a; E) = \frac{i\hbar}{2Mc} \int_0^\infty dL \int \mathcal{D}\rho(\lambda) \Phi[\rho(\lambda)] \times \int_{x_a}^{x_b} \mathcal{D}x(\lambda) \exp \left\{ -\frac{1}{\hbar} \int_{\lambda_a}^{\lambda_b} d\lambda \left[\frac{M}{2\rho(\lambda)} x'^2(\lambda) - \rho(\lambda) \frac{(E - V(x))^2}{2Mc^2} + \rho(\lambda) \frac{Mc^2}{2} \right] \right\} \quad (19)$$

$$= \frac{\begin{vmatrix} G_0(x_b, x_a; E) & G_0(x_b, b; E) & G_0(x_b, a; E) \\ G_0(b, x_a; E) & G_0(b, b; E) & G_0(b, a; E) \\ G_0(a, x_a; E) & G_0(a, b; E) & G_0(a, a; E) \end{vmatrix}}{\begin{vmatrix} G_0(b, b; E) & G_0(b, a; E) \\ G_0(a, b; E) & G_0(a, a; E) \end{vmatrix}}, \quad (20)$$

where the notation G_0 stands for the non-perturbative relativistic fixed-energy amplitude.

The methods present in this paper enable us to solve a large class of the relativistic potential problems. It is our hope that our studies would help to achieve the ultimate goal of obtaining a comprehensive and complete path integral description of quantum mechanics and quantum field theory, including quantum gravity and cosmology.

Acknowledgments

D-H Lin was supported by the National Youth Council of the ROC under contract number NYC300375. Part of the work is supported by the rotary club of Hsin Chu north west district.

References

1. H. Kleinert, “*Path Integrals in Quantum Mechanics, Statistics and Polymer Physics*,” World Scientific, Singapore, 1995.
2. D. H. Lin, J. Phys. **A** **30**, 3201 (1997).
3. D. H. Lin, J. Phys. **A** **30**, 4365 (1997).
4. D. H. Lin, J. Phys. **A** **31**, 4785 (1998).
5. D. H. Lin, hep-th/9708144; hep-th/9709152.